arXiv:hep-ex/0411087v1 29 Nov 2004

Hermes Precision Results on g_1^{p}, g_1^{d} and g_1^{n} and the First Measurement of the Tensor Structure Function b_1^{d}

Caroline Riedl *(on behalf of the* HERMES *collaboration)*
Physikalisches Institut 2 der Universität Erlangen-Nürnberg,
Erwin-Rommel-Str. 1, D-91058 Erlangen, Germany
E-mail: criedl@mail.desy.de

Abstract. Final HERMES results on the proton, deuteron and neutron structure function g_1 are presented in the kinematic range $0.0021 < x < 0.9$ and $0.1 < Q^2 < 20$ GeV^2. These results base on a refined analysis and are corrected for radiative and detector smearing effects using an unfolding algorithm. Furthermore, preliminary results on the first measurement of the tensor asymmetry A_{zz} and the tensor structure function b_1^{d} are presented.

Over the last 15 years, measurements of the polarized structure function g_1 by inclusive polarized deep-inelastic scattering (DIS)[1] have shed light on one basic ingredient of the spin puzzle: the quark contribution to the nucleon spin, $\Delta\Sigma$. For a spin-1 target, besides g_1 the quadrupole structure function b_1 is needed to parameterize the hadronic part of the interaction.[2]

The final HERMES g_1 results presented here base on a refined analysis of data taken with longitudinally polarized hydrogen[3] and deuterium[4] targets (4 resp. $11.5 \cdot 10^6$ DIS events), the b_1 results on a separate data set with a tensor-polarized deuterium target ($2.4 \cdot 10^6$ DIS events).[5] The kinematic range covers $0.0021 < x < 0.9$ and $0.1 < Q^2 < 20$ GeV^2.

In the fixed-target experiment HERMES, the 27.6 GeV lepton beam of the HERA storage ring is scattered off an internal target of pure gases. The lepton beam is transversely self-polarized, for the reported data reaching an average polarization of 53%; longitudinal beam polarization in the interaction region is obtained by pairs of spin rotators. The HERMES atomic gas target[6] consists of an Atomic Beam Source (ABS) to generate and a Breit Rabi Polarimeter to measure nuclear polarization; a tubular open-ended, cooled storage cell is mounted inside the beam pipe confining the gas fed from the ABS. The average vector polarization was 85%, furthermore, the target design allowed for a simultaneous setting of an average tensor polarization of −166% and a residual vector polarization of only +1%, which makes it possible to decouple the b_1 from the g_1 measurement.

The HERMES forward spectrometer[7] includes numerous tracking chambers in front and behind a 1.3 Tm magnetic field and various detectors for particle identification. Electrons and hadrons are discriminated using a likelihood method based on the combination of four detectors.[8] The presented inclusive sample is contaminated less than 1% with hadrons.

The spin structure function $g_1(x, Q^2)$ is for both the proton and the deuteron determined from the ratio g_1/F_1 which is approximately equal to the virtual photon asymmetry A_1 measured via the longitudinal cross section asymmetry $A_{\|}$. For analysis details, see e.g. Ref. [3]. For the neutron, g_1^{n} is calculated as a linear combination of g_1^{p} and g_1^{d}.

The measured asymmetries have been corrected for detector smearing and QED radiative effects to obtain the Born asymmetries.[8,9] These corrections have been applied using an unfolding algorithm that keeps track of the kinematic migration of events and doesn't require a fit on the data. The final Born asymmetries depend only on the measured data, on the detector model, on the known unpolarized cross sections and on the model for background processes. Figure 1 shows the smearing unfolded g_1/F_1 and xg_1 results and, based on the simulated event migration, their statistical correlations which must be taken into account when e.g. calculating errors on moments or performing QCD fits to g_1 data.[9] The applied algorithm removes systematic correlations between data points. A compilation of world data on g_1/F_1 and xg_1 is shown in Fig. 2.[10,11,12]

The tensor structure function $b_1(x, Q^2)$ is obtained from the ratio b_1/F_1, measured via A_{zz} which compares the helicity-0 state of the deuteron with its averaged non-zero states. In Fig. 3, A_{zz} and b_1 are displayed as measured for the first time by HERMES; A_{zz} is only of the order of 1%, implying the quadrupole contribution to the measurement of g_1 to be negligible. The steep rise of b_1 for small-x suits latest model calculations.[5] The unfolding algorithm discussed above will also be applied to these data.[13]

Acknowledgments

I thank my HERMES colleagues, especially the inclusive group. Furthermore, I gratefully acknowledge the financial support by the German Bundesministerium für Bildung und Forschung (contract number 06 ER 125I).

References

1. For a compilation of polarized data see: `http://durpdg.dur.ac.uk/HEPDATA`
2. P. Hoodbhoy *et al.*, Nucl. Phys. B**312** (1989) 571



Figure 1. Smearing unfolded g_1/F_1 (left) and xg_1 (middle) as measured by HERMES and the correlation matrix between statistical errors (right), for the proton (top row) and for the deuteron (bottom row). Error bars are statistical, displaying only the diagonal element of the correlation matrix, the shaded bands show the estimated systematic uncertainties

3. HERMES Collaboration, A. Airapetian *et al.*, Phys. Lett. B**442** (1998) 484
4. L. De Nardo, for the HERMES Collaboration, *Proceedings from PRAHA-Spin-2001, Prague, Czech Republic, Jul 15 - 28, 2001*
5. M. Contalbrigo, for the HERMES Collaboration, *Proceedings from SPIN 2002, New York, USA, Sep 9 - 14, 2002*, hep-ex/0211014
6. HERMES Collaboration, A. Airapetian *et al.*, Nucl. Instr. and Meth. A (in press), hep-ex/0408013
7. HERMES Collaboration, K. Ackerstaff *et al.*, Nucl. Instr. A**417** (1998) 230
8. HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. D (in press), hep-ex/0407032
9. HERMES Collaboration, publication on g_1 in preparation
10. SMC Collaboration, B. Adeva *et al.*, Phys. Rev. D**60** (1999) 072004
11. E143 Collaboration, K. Abe *et al.*, Phys. Rev. D**58** (1998) 112003
12. E155 Collaboration, P.L. Anthony *et al.*, Phys. Lett. B**463** (1999) 339



Figure 2. HERMES g_1/F_1 (left) and xg_1 data (right) in comparison to SLAC and CERN data for the proton (top panels) and the deuteron (middle panels) and xg_1 for the neutron from p and d data (right bottom panel) at the measured Q^2 (left bottom). Error bars show the quadratic sum of statistical and systematic uncertainties.



Figure 3. The HERMES tensor asymmetry A_{zz} (left) and tensor structure function b_1^{d} (right; lower panel: average Q^2 for the measurements). The error bars are statistical only, the shaded bands show the estimated systematic uncertainties.

13. HERMES Collaboration, publication on b_1 in preparation